For Chengdu Workshop on ASIAEX 2001

Characteristic of the Internal Waves at Location of 126°54.32'E/ 29°24.01'N

Shihong ZHOU

(Dalian Scientific Test & Control Technology Institute, China)

Jixun ZHOU Jie YANG

(Georgia Institute of Technology, Atlanta, USA)

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to completing and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar DMB control number.	ion of information. Send comments arters Services, Directorate for Information	regarding this burden estimate of mation Operations and Reports	or any other aspect of th , 1215 Jefferson Davis I	is collection of information, Highway, Suite 1204, Arlington		
1. REPORT DATE 01 DEC 2002		2. REPORT TYPE N/A		3. DATES COVERED			
4. TITLE AND SUBTITLE				5a. CONTRACT	NUMBER		
	he Internal Waves a	t Location of		5b. GRANT NUM	1BER		
126degree54.32'E/29degree24.01'N					5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)					5d. PROJECT NUMBER		
	5e. TASK NUMBER						
					5f. WORK UNIT NUMBER		
7. PERFORMING ORGANI Dalian Scientific Technol Institute of Technol	8. PERFORMING ORGANIZATION REPORT NUMBER						
9. SPONSORING/MONITO	10. SPONSOR/MONITOR'S ACRONYM(S)						
	11. SPONSOR/MONITOR'S REPORT NUMBER(S)						
12. DISTRIBUTION/AVAII Approved for publ	LABILITY STATEMENT ic release, distributi	on unlimited					
13. SUPPLEMENTARY NO Also see: M001452	otes , The original docum	nent contains color	images.				
14. ABSTRACT							
15. SUBJECT TERMS							
16. SECURITY CLASSIFIC	17. LIMITATION OF	18. NUMBER	19a. NAME OF				
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	ABSTRACT OF PAGES RI UU 20		RESPONSIBLE PERSON		

Report Documentation Page

Form Approved OMB No. 0704-0188

What the topic focuses on?

- □Observation of Internal Waves in ECS ASIAEX 2001
- ☐ Features of the Temperature Fluctuation
- □ Characteristic of the Vertical Displacement Power Spectrum of Internal Waves
- □Normal mode of internal wave and its dispersion including Phase velocities, group velocities, and eigenfunctions of normal modes

1. Introduction

Experiment Site:

126°54.32'E/29°24.01'N

Water depth of sea area: ~ 105m

2. Observation of Internal Waves

Survey Duration:

~ 63 hours from 08:40/Jun.3 to 0:00/Jun.6/2001

Survey Instruments:

17 WaDar Temperature Sensors from GIT

Sensor depths:

19.95, 23.1, 26.25, 29.4, 32.55, 35.7, 38.85, 42, 45.15,

48.3, 51.45, 54.6, 57.75, 60.9, 65.1, 71.4 and 79.8m




Figure 1 Sea water temperature data curves at different depth from 08:40/Jun.3 to 00:00/Jun.6/2001

Features of the Temperature Fluctuation:

- ☐ Three elevation solitons at 53:15, 53:46 and 54:14 which are separated from internal tides
- ☐ Thermocline Layer: from 20m to 75m
- ☐ Internal semi-diurnal tides with strong fluctuations at 8:00, 21:00, 33:00, 46:00 and 58:00
- □Cool water-mass between 11:48 and 13:52




Figure 2 Sea water temperature distribution plot at different depth from 08:40/Jun.3 to 00:00/Jun.6/2001

3. Characteristic of the Vertical Displacement Power Spectrum of Internal Waves





Figure 3 Vertical displacement curves at different temperature from 08:40/Jun.3 to 00:00/Jun.6/2001






Figure 4 Power spectrum of vertical particle displacement at different temperature

Table 1 The attenuation coefficients of vertical displacement power spectrum at different temperature

T(°C)	19	19.5	20	20.5	21	21.5	22
b	1.9065	1.4571	1.503	1.4803	1.4923	1.6636	1.3848

During the frequency range of $w_i < w < N(z)$, the spectrum attenuation coefficients of the vertical displacement power spectrum are about -1.5

- □ Inertial frequency: $w_i = 2Ω \sin j = 0.0043$ cpm
- **Brunt-Vaisala's frequency:** N(z)

4. Normal mode of internal wave and its dispersion

Wave Equation of internal wave:

$$\frac{\partial^2 \mathbf{y}_j(z)}{\partial z^2} + \left[\frac{N^2(z) - \mathbf{w}^2}{\mathbf{w}^2 - \mathbf{w}_i^2} \right] k_{hj}^2 \mathbf{y}_j(z) = 0$$

 k_{hi} : horizontal wave-number of j^{th} normal mode of internal wave

Brunt-Vaisala's frequency:

$$N^{2}(z) = -\frac{g}{\mathbf{r}(z)} \frac{\partial \mathbf{r}(z)}{\partial z} = ag \frac{\partial T(z)}{\partial z} - bg \frac{\partial S(z)}{\partial z}$$


Eigenfunction Approximation by Talor's Series Expansion:

$$\mathbf{y}_{j}^{n+1} = \mathbf{y}_{j}^{n} + \mathbf{y}_{j}^{n} h + \mathbf{y}_{j}^{n} \frac{h^{2}}{2!} + \mathbf{y}_{j}^{n} \frac{h^{3}}{3!} + \cdots$$

$$\mathbf{y}_{j}^{n-1} \approx \mathbf{y}_{j}^{n} - \mathbf{y}_{j}^{n} h + \mathbf{y}_{j}^{n} \frac{h^{2}}{2!} - \mathbf{y}_{j}^{n} \frac{h^{3}}{3!} + \cdots$$

$$\mathbf{y}_{j}^{n} \approx \frac{\mathbf{y}_{j}^{n+1} - \mathbf{y}_{j}^{n}}{h} + \left[\frac{N^{2}(z_{n}) - \mathbf{w}^{2}}{\mathbf{w}^{2} - \mathbf{w}_{i}^{2}} \right] k_{hj}^{2} \mathbf{y}_{j}^{n} \frac{h}{2}$$

$$\mathbf{y}_{j}^{n} \approx \frac{\mathbf{y}_{j}^{n} - \mathbf{y}_{j}^{n-1}}{h} - \left[\frac{N^{2}(z_{n}) - \mathbf{w}^{2}}{\mathbf{w}^{2} - \mathbf{w}_{i}^{2}} \right] k_{hj}^{2} \mathbf{y}_{j}^{n} \frac{h}{2}$$



$$\mathbf{y}_{j}^{n-1} + \left\{ -2 + h^{2} \left[\frac{N^{2}(z_{n}) - \mathbf{w}^{2}}{\mathbf{w}^{2} - \mathbf{w}_{i}^{2}} \right] k_{hj}^{2} \right\} \mathbf{y}_{j}^{n} + \mathbf{y}_{j}^{n+1} = 0, \qquad n = 1, 2, \dots, N-1$$

Boundary Condition:

$$\mathbf{y}_{j}^{0} = 0$$
 and $\mathbf{y}_{j}^{N} = 0$

Wave Equation:

$$\mathbf{A}(k_{hj}^2)$$
? $_j = 0 \implies$ Eigenfunctions

Eigenvalue Equation:

$$\det[\mathbf{A}(k_{hj}^2)] = 0 \implies \frac{\text{Phase/Group Velocities}}{\text{of Normal Modes}}$$




Figure 5 Phase velocities, group velocities and eigenfunctions of the first five normal modes of internal wave

At frequency of 0.05cpm, Phase/Group Velocities:

 \square 1th normal mode: 1.77m/s and 1.73m/s

□2th normal mode: 0.73m/s and 0.71m/s

□3th normal mode: 0.44m/s and 0.43m/s

□4th normal mode: 0.34m/s and 0.33m/s

□5th normal mode: 0.27m/s and 0.26m/s

?
$$_{j} = [\mathbf{y}_{j}^{0}, \mathbf{y}_{j}^{1}, \mathbf{y}_{j}^{2}, \cdots, \mathbf{y}_{j}^{N}]$$

$$\mathbf{A}\mathfrak{L}^{1} = \begin{bmatrix} 1 & 0 & & & & & & \\ 1 & g_{1} & 1 & & & & & \\ & 1 & g_{2} & 1 & & & & \\ & & \ddots & \ddots & \ddots & & \\ & & & 1 & g_{N-2} & 1 & \\ & & & & 1 & g_{N-1} & 1 \\ & & & & & 0 & 1 \end{bmatrix}_{(N+1)\times(N+1)}$$

Here,
$$g_n = -2 + h^2 \left[\frac{N^2(z_n) - \mathbf{w}^2}{\mathbf{w}^2 - \mathbf{w}_i^2} \right] k_{hj}^2$$
 $n = 1, 2, \dots, N-1$

5. Conclusion

- (1) The internal wave owns the feature of internal semidiurnal tides with strong fluctuations.
- (2) There exist three elevation solitons which are separated from internal tides.
- (3) The mean spectrum attenuation coefficient of the vertical displacement power spectrum is about -1.5.
- (4)The estimated mean group velocities of five normal modes of internal wave at frequency of 0.05cpm are 1.73m/s, 0.71m/s, 0.43m/s, 0.33m/s and 0.26m/s, respectively.

THANKS